arXiv:1402.6981v1 [math.NA] 27 Feb 2014

Integrators on homogeneous spaces: Isotropy choice and connections

Hans Munthe-Kaas*[1] and Olivier Verdier†[2]

[1] *Department of Mathematics, University of Bergen, Norway*
[2] *Department of Mathematics and Statistics, Ume[Pleaseinsert"PrerenderUnicode–"intopreamble] University, Sweden*

2014-02-28

We consider numerical integrators of ODEs on homogeneous spaces (spheres, affine spaces, hyperbolic spaces). Homogeneous spaces are equipped a built-in symmetry. A numerical integrator respects this symmetry if it is equivariant. One obtain homogeneous space integrators by combining a Lie group integrator with an isotropy choice. We show that equivariant isotropy choices combined with equivariant Lie group integrators produces an equivariant homogeneous space integrator. Moreover, we show that the RKMK, Crouch–Grossman or commutator-free methods are equivariant. In order to show this, we give a novel description of Lie group integrators in terms of stage trees and motion maps, which unifies the known Lie group integrators. We then proceed to study the equivariant isotropy maps of order zero, which we call connections, and show that they can be identified with reductive structures and invariant principal connections. We give concrete formulas for connections in standard homogeneous spaces of interest, such as Stiefel, Grassmannian, isospectral, and polar decomposition manifolds. Finally, we show that the space of matrices of fixed rank possesses no connection.

Keywords Homogeneous spaces · Symmetric spaces · Lie Group integrators · Connection · Runge–Kutta · Skeleton · Stiefel Manifold · Lax Pair · Grassmannian · Projective space · Polar decomposition · Constant rank matrices

Mathematics Subject Classification (2010) 22F30 · 53C30 · 14M15 · 65Pxx · 37Mxx

*hans.munthe-kaas@math.uib.no
†olivier.verdier@math.umu.se

Contents

1 Introduction **3**

2 Isotropy Maps **6**
- 2.1 Equivariance . . . 6
- 2.2 Locality . . . 7
- 2.3 Fundamental Example: the tautological isotropy map . . . 7

3 Runge–Kutta Methods **8**
- 3.1 Skeletons . . . 8
- 3.2 Transition Functions . . . 9
- 3.3 Examples . . . 11
  - 3.3.1 Runge–Kutta methods in $\mathbf{R}^d$ . . . 12
  - 3.3.2 RKMK4 . . . 12
  - 3.3.3 CG3 . . . 13
  - 3.3.4 Symmetric Gauss of order four . . . 13
- 3.4 Order Zero and Exact Solution . . . 14
- 3.5 Equivariance of Skeletons . . . 14
- 3.6 Locality . . . 15

4 Connections and Reductivity **15**
- 4.1 Connections . . . 15
- 4.2 Reductive decompositions . . . 18
- 4.3 Relation with Invariant Principal Connections . . . 19
- 4.4 Flatness and Symmetry . . . 21
- 4.5 Existence of Connections . . . 21
- 4.6 Descending Property of Skeletons . . . 22

5 Connection Examples **22**
- 5.1 Lie Group as Homogeneous Spaces . . . 22
  - 5.1.1 Left and Right Actions . . . 22
  - 5.1.2 Cartan–Schouten Action . . . 24
- 5.2 Affine spaces . . . 25
- 5.3 Stiefel Manifolds and spheres . . . 26
- 5.4 Isospectral Flows, Lax Pairs, Grassmannians and Projective Spaces . . . 28
  - 5.4.1 The symmetric case: Grassmannians . . . 31
  - 5.4.2 The flat case: integrable Lax pairs . . . 31
- 5.5 Polar Decompositions . . . 32
- 5.6 Matrices of fixed rank . . . 32

6 Conclusion and Open Problems **34**

1 Introduction

The purpose of this paper is to give concrete formulas and definitions for integrators on *homogeneous spaces*. A homogeneous space is a manifold $\mathcal{M}$ on which a group G acts transitively. Homogeneous spaces are ubiquitous in numerical analysis and computer vision. Examples include spheres, Stiefel manifolds and Grassmannians [9], symmetric positive definite matrices used in diffusion tensor computing [20], isospectral manifolds used in the context of Lax pairs [3], and constant rank matrices used in low-rank approximation [11, § IV.9.3]. We study all these cases in thorough details in § 5.

The most convenient way to construct an integrator on a homogeneous space is, as noted in [13] to use a Lie group integrator on G along with an *isotropy map*.

There is, however, an essential ambiguity in the isotropy map, as many infinitesimal motions correspond to a given vector field, as illustrated in Figure 1. One key idea to address this issue is the notion of *equivariance*, which was already developed in [17]. Indeed, a homogeneous space has a built-in symmetry which is only preserved if the integrator at hand is equivariant. We will show in Proposition 3.2 that if the isotropy map is equivariant, then so is the corresponding integrator. This already drastically limits the choices of isotropy maps.

A further limitation to the choice of isotropy map is *locality*, also introduced in a similar guise in [17]. The simplest possible local isotropy maps are isotropy *forms*, i.e., $\mathfrak{g}$-values one-forms, which we regard as order zero isotropy maps. In the remaining of the paper, we focus on the equivariant isotropy maps of order zero, which we call *connections*, as they can be identified with Nomizu's invariant principal connections and reductive structures, see Figure 4.

We then thoroughly examine existence, and concrete formulas, for connections on homogeneous spaces that have been used in numerical analysis, computer visions and optimisation. The reader will find a summary of the results in Table 1.

Besides the definitions of equivariant and local isotropy maps in § 2, and the definition of connections in § 4, the following results appearing in this paper are new:

- Proof of the equivalence of connections (order zero equivariant isotropy maps), reductive structures and principal invariant connections in § 4
- Recognising that the formulas in Table 1, when they existed, are in fact connections
- Showing in Proposition 5.7 that the homogeneous space of fixed rank matrices has no connection.
- Giving in § 3 an encompassing definition of the existing Lie group methods, and showing that they are equivariant (Proposition 3.2) and local (Proposition 3.5).
- Proof that if the isotropy choice must be equivariant, then so is the corresponding method (Proposition 3.2).

We now recall some basic definitions about homogeneous spaces that will be needed throughout this paper, and refer to [22], [15] and [10] for a thorough exposition. The



(a) Motion ambiguity



(b) Infinitesimal motion ambiguity

Figure 1: We illustrate the concept of a connection in the case of the Euclidean displacement group $\mathsf{SO}(2) \ltimes \mathbf{R}^2$ acting on $\mathbf{R}^2$.
(a) There are plenty of displacements bringing the point x_0 to the point x_1. The ambiguity in the displacement choice is measured by the isotropy group $\mathsf{SO}(2)$, which, in this case, has dimension one. The displacements consist of either a translation, or arcs of circle with centre located on the perpendicular bisector of x_0 and x_1. The discovery of Nomizu [18] is that there is one *preferred* choice: the translations, marked in bold in the picture. The reason translations stand apart, is that they are geodesics associated to an invariant connection. In our framework, they are associated to a connection.
(b) Suppose that a choice was made to use circular motions, say, on the right, which gives the point $\exp(\xi) \cdot x$ on the picture. As the isotropy choice is a linear map, the isotropy choice of the opposite of this vector commands to follow $\exp(-\xi) \cdot x$, i.e., to turn *left*. On the other hand, rotating the initial vector by a half turn would command to turn *right*. This choice discrepancy illustrates the lack of equivariance. In Proposition 5.1, we show rigorously that translations are the only equivariant choices for full affine spaces corresponding to $\mathsf{GL}(d) \ltimes \mathbf{R}^d$ acting on $\mathbf{R}^d$.



Figure 2: An illustration of a homogeneous space. The origin $\mathsf{o} \in \mathcal{M}$ is an arbitrary point of $\mathcal{M}$ which defines the corresponding isotropy group H. Any point $x \in \mathcal{M}$ may then be identified with the set $\bar{g}\,\mathsf{H}$, for some $\bar{g}$ such that $x = [\bar{g}] = \bar{g}\,\mathsf{H}$. Multiplication on the right by an element $h \in \mathsf{H}$ projects to the same point, i.e., $x = [\bar{g}h] = [\bar{g}]$. Note that the multiplication on the left by an element $g \in \mathsf{G}$ sends fibres to fibres, and descends to $\mathcal{M}$ to the action of g to $\mathcal{M}$, i.e., $[g\,\bar{g}] = g \cdot [\bar{g}]$.

reader will find a summary of the concepts and definitions of homogeneous spaces in Figure 2.

We use the standard notations pertaining to calculus on smooth manifolds [1]. Given a manifold $\mathcal{M}$, we denote its tangent bundle by $\mathsf{T}\mathcal{M}$. The space of sections of that tangent bundle (the vector fields) is denoted by $\mathfrak{X}(\mathcal{M})$. Given a function $f\colon \mathcal{M} \to \mathcal{N}$, where $\mathcal{M}$ and $\mathcal{N}$ are two smooth manifolds, we denote its push-forward, or Jacobian, by $\mathsf{T}f$, so $\mathsf{T}f\colon \mathsf{T}\mathcal{M} \to \mathsf{T}\mathcal{N}$.

A ***homogeneous space*** is a smooth manifold $\mathcal{M}$, equipped with a transitive action of a Lie group G [15, 14, 22, 10]. We denote this transitive action as

$$g \cdot x, \qquad g \in \mathsf{G} \quad x \in \mathcal{M}. \tag{1}$$

Fixing an arbitray point $\mathsf{o} \in \mathcal{M}$, we define the ***isotropy group*** at this point by

$$\mathsf{H} \coloneqq \{\, g \in \mathsf{G} \mid g \cdot \mathsf{o} = \mathsf{o} \,\}. \tag{2}$$

The manifold $\mathcal{M}$ may be identified with cosets

$$x \equiv g\,\mathsf{H} = \{\, g\,h \mid h \in \mathsf{H} \,\}. \tag{3}$$

We denote the projection from G to $\mathcal{M}$ by

$$[g] \coloneqq g\,\mathsf{H}. \tag{4}$$

For a tangent vector X at a point $g \in \mathsf{G}$, we denote the projection by an abuse of notation

$$[X] \coloneqq \mathsf{T}\pi\, X, \tag{5}$$

where π denotes the projection $\pi(g) = [g]$.

Note that in order to simplify the notations, we will always assume that G is a subgroup of $\mathsf{GL}(d)$ for some integer d. In particular, we will consider the ***Lie algebra*** $\mathfrak{g}$, the tangent space at the identity of G, to be an affine subspace of $\mathbf{R}^{d\times d}$, the space of $d \times d$ matrices. This allows us to define operations such as the adjoint action of $g \in \mathsf{G}$ on $\xi \in \mathfrak{g}$ by matrix multiplication as $\xi \to g\,\xi\,g^{-1}$. Similarly, the multiplication $g\,\xi$ denotes the tangent of the left multiplication map applied to $\xi \in \mathfrak{g}$, and is thus considered as a vector at g, i.e., $g\,\xi \in \mathsf{T}_g\mathsf{G}$.

Finally, will repeatedly use that the space $\mathsf{T}_{\circ}\mathcal{M}$ is isomorphic to $\mathfrak{g}/\mathfrak{h}$, which is straightforward to check [22, § 4.5].

2 Isotropy Maps

We define isotropy maps and give some definitions, especially the notion of equivariance.

2.1 Equivariance

Definition 2.1. We call an ***isotropy map*** a linear map

$$\varpi \in \mathcal{L}\big(\mathfrak{X}(\mathcal{M}), \mathcal{C}^\infty(\mathcal{M}, \mathfrak{g})\big) \tag{6}$$

which satisfies the ***consistency condition***

$$\langle \varpi, f\rangle(x) \cdot x = f(x) \qquad x \in \mathcal{M} \quad f \in \mathfrak{X}(\mathcal{M}). \tag{7}$$

The interpretation is that, given a vector field f, the corresponding value $\nu = \langle \varpi, f\rangle \in \mathcal{C}^\infty(\mathcal{M}, \mathfrak{g})$ associates to every point $x \in \mathcal{M}$ a *frozen vector field* F^x based at x defined by the infinitesimal action

$$F^x(y) = \nu(x) \cdot y \qquad y \in \mathcal{M}. \tag{8}$$

The consistency condition (7) means that the frozen vector field for f based at x, and evaluated at x, coincides with f at x, i.e., $F^x(x) = f(x)$.

Definition 2.2. An isotropy map is ***equivariant*** if

$$\langle \varpi, g \cdot f\rangle = g \cdot \langle \varpi, f\rangle \tag{9}$$

where the action of $g \in \mathsf{G}$ on a vector field f is given by [14, § 2.4]

$$(g \cdot f)(x) \coloneqq \mathsf{T} g \, f(g^{-1} \cdot x), \qquad x \in \mathcal{M}, \tag{10}$$

the action on $\nu \in \mathcal{C}^\infty(\mathcal{M}, \mathfrak{g})$ is

$$(g \cdot \nu)(x) \coloneqq g \cdot \nu(g^{-1} \cdot x), \qquad x \in \mathcal{M}, \tag{11}$$

and the action on $\xi \in \mathfrak{g}$ is the *adjoint action* [14, § 2.6]

$$g \cdot \xi = g \, \xi \, g^{-1}. \tag{12}$$

Putting all the bits of the definitions together, equivariance of the isotropy map ϖ is written as

$$\langle \varpi, g \cdot f \rangle(x) = g \, \langle \varpi, f \rangle(g^{-1} \cdot x) \, g^{-1}, \qquad x \in \mathcal{M}, \quad g \in \mathsf{G}. \tag{13}$$

2.2 Locality

Recall that the *support* $\operatorname{supp}(\nu)$ of a section ν of a vector bundle is the closure of the set where the section is non-zero. For an isotropy choice $\nu \in \mathcal{C}^\infty(\mathcal{M}, \mathfrak{g})$, this gives

$$\operatorname{supp}(\nu) \coloneqq \overline{\{\, x \in \mathcal{M} \mid \nu(x) \neq 0 \,\}}, \tag{14}$$

and for a vector field $f \in \mathfrak{X}(\mathcal{M})$,

$$\operatorname{supp}(f) \coloneqq \overline{\{\, x \in \mathcal{M} \mid f(x) \neq 0 \,\}}. \tag{15}$$

Definition 2.3. An isotropy map $\varpi \in \mathcal{L}\big(\mathfrak{X}(\mathcal{M}), \mathcal{C}^\infty(\mathcal{M}, \mathfrak{g})\big)$ is ***local*** if it is support non increasing, i.e.,

$$\operatorname{supp}\big(\langle \varpi, f \rangle\big) \subset \operatorname{supp}(f) \qquad f \in \mathfrak{X}(\mathcal{M}). \tag{16}$$

2.3 Fundamental Example: the tautological isotropy map

A fundamental example, on any manifold $\mathcal{M}$, is given by the action of the *whole group of diffeomorphisms* $\mathsf{G} = \mathsf{Diff}(\mathcal{M})$. The action is transitive. Note that the Lie algebra $\mathfrak{g}$ is now $\mathfrak{g} = \mathfrak{X}(\mathcal{M})$, where $\mathfrak{X}(\mathcal{M})$ is the space of vector fields on $\mathcal{M}$ regarded as a Lie algebra.

We define the ***tautological isotropy map*** [1] by

$$\langle \varpi, f \rangle(x) \coloneqq f, \qquad \forall x \in \mathcal{M}, \quad f \in \mathfrak{X}(\mathcal{M}). \tag{17}$$

The tautological isotropy map is an *equivariant* but *non local* isotropy map.

Indeed, $\langle \varpi, f \rangle$ is a *constant function* from $\mathcal{M}$ to $\mathfrak{g} = \mathfrak{X}(\mathcal{M})$. The support of a constant function is either $\mathcal{M}$, or, if the constant is zero, the empty set. This shows that ϖ is *not local*, as the support of a non-zero vector field need not be the whole manifold $\mathcal{M}$.

The reason this isotropy map is fundamental is that, as we shall see in § 3.4, it is the isotropy map that, assuming zero-order of the underlying skeleton, gives the exact solution.

[1] With the notation (49) that isotropy choice is simply $\langle \varpi, f \rangle = f$ hence the name of *tautological* isotropy map.



Figure 3: A summary of the relation between the constitutive elements of a method on a homogeneous space.

3 Runge–Kutta Methods

We put the known Lie–Group and homogeneous space methods in a unified framework. This allows us to show in Proposition 3.2 that if isotropy map is equivariant, then so is the corresponding integrator.

3.1 Skeletons

We define a Runge–Kutta skeleton for a given Lie group G from the following ingredients.

Stage Tree Γ Tree (i.e., a connected, undirected graph with no cycles), in which two vertices are singled out: the ***initial vertex***, denoted "$\circ$" and the ***final vertex*** denoted "$\bullet$".

Motions $\Psi_{i,j}$ Maps

$$\Psi_{i,j} \colon \mathfrak{g}^n \to \mathsf{G} \tag{18}$$

defined for any two adjacent vertices i, j in the graph, and the compatibility

$$\Psi_{i,j} = \Psi_{j,i}^{-1}. \tag{19}$$

One defines a skeleton

$$\Sigma \colon \mathcal{C}^\infty(\mathcal{M}, \mathfrak{g}) \to \mathsf{Diff}(\mathcal{M}) \tag{20}$$

using the *stages* $X_i \in \mathcal{M}$, and the *frozen vector fields* $F_i \in \mathfrak{g}$ as intermediate variables.

From a skeleton and an isotropy map, we obtain an integrator on any homogeneous manifold $\mathcal{M}$ for which G is the symmetry group. Indeed, given an isotropy choice $\nu \in \mathcal{C}^\infty(\mathcal{M}, \mathfrak{g})$, the map

$$\Sigma(\nu) \in \mathsf{Diff}(\mathcal{M}) \tag{21}$$

is defined by

$$x_1 = \Sigma(\nu)(x_0) \tag{22}$$

as follows.

Runge–Kutta Method

$$X_\circ = x_0 \tag{23a}$$
$$X_i = \Psi_{i,j}(F) \cdot X_j \qquad \forall \text{ edge } i,j \tag{23b}$$
$$F_i = \nu(X_i) \qquad \forall \text{ vertex } i \tag{23c}$$
$$x_1 = X_\bullet \tag{23d}$$

In the sequel we will always assume that ν is scaled (with a time step), so that the equations (23) implicitly defining x_1 have exactly one solution.

Let us consider the simplest possible case, a stage tree containing only the initial vertex $\circ$ and final vertex $\bullet$:

(24)

The corresponding equations for an isotropy choice $\nu = \langle \varpi, f \rangle$ corresponding to an isotropy map ϖ, are

$$X_\bullet = \Psi_{\bullet,\circ}(F_\circ, F_\bullet) \cdot X_\circ, \tag{25a}$$
$$F_\circ = \langle \varpi, f \rangle(X_\circ), \tag{25b}$$
$$F_\bullet = \langle \varpi, f \rangle(X_\bullet), \tag{25c}$$

and the integrator maps the initial condition $X_\circ$ to the point $X_\bullet$.

Possible values for $\Psi_{\bullet,\circ}(F_\circ, F_\bullet)$ are $\exp(F_\circ)$, which gives the forward Euler method, $\exp(F_\bullet)$, which gives the backward Euler method, or $\exp\big((F_\circ + F_\bullet)/2\big)$ which gives the trapezoidal rule.

3.2 Transition Functions

Note that most often, the motions $\Psi_{i,j}$ are defined by

$$\Psi_{i,j} \coloneqq \Psi \circ \theta_{i,j} \tag{26}$$

with

Motion map Ψ

$$\Psi\colon \mathfrak{g} \to \mathsf{G}, \tag{27}$$

which is usually either the exponential map, or an approximation of it, or the Cayley map when G is quadratic; it should have the property that

$$\Psi(-\xi) = \Psi(\xi)^{-1}, \tag{28}$$

which holds both for the exponential and Cayley map.

Transition functions $\theta_{i,j}$ For any edge i, j, the transition function

$$\theta_{i,j}\colon \mathfrak{g}^n \to \mathfrak{g}, \tag{29}$$

with compatibility

$$\theta_{i,j}(F) = -\theta_{j,i}(F). \tag{30}$$

We will give the full expression of $\theta_{i,j}$ on the stage tree as



(31)

For example, the *forward Euler method* is given by

$$\theta_{\bullet,\circ} = F_\circ \tag{32}$$

so this is equivalent to the tree



(33)

Note that (30) means that the orientation is arbitrary, so with a different orientation, we obtain exactly *the same method*, as $\theta_{\circ,\bullet} = -\theta_{\bullet,\circ}$:



(34)

We shade the vertices which value is not used in any of the transition functions. In the case above, the value $F_\bullet$ is never needed, so the vertex $\bullet$ is shaded.

With these notations, the *backward Euler method* is given by the transition function $\theta_{\bullet,\circ} = F_\bullet$, so we write

(35)

For the trapezoidal rule, the transition function is $\theta_{\bullet,\circ} = (F_\circ + F_\bullet)/2$, so it is written as:



(36)

A slightly more involved example is the implicit midpoint rule, which requires an extra stage which has the arbitrary label "$\star$". The stage tree has thus three vertices, $\circ$, $\bullet$ and $\star$. The transition functions are

$$\theta_{\star,\circ} = F_\star/2 \tag{37a}$$
$$\theta_{\bullet,\circ} = F_\star \tag{37b}$$

so we write



(38)

Note again that the values $F_\circ$ and $F_\bullet$ are not used, so we shade the corresponding vertices $\circ$ and $\bullet$. We also emphasise the (unique) path between the initial and final vertices.

For the sake of completeness, we also give the corresponding equations, which are

$$X_\star = \exp(F_\star/2) \cdot X_\circ \tag{39a}$$
$$X_\bullet = \exp(F_\star) \cdot X_\circ \tag{39b}$$

and

$$F_\star = \nu(f)(X_\star) \tag{40}$$

According to our convention, the integrator maps $X_\circ$ to $X_\bullet$. Note that $X_\circ$, $X_\bullet$ and $X_\star$ are points in $\mathcal{M}$, and that $F_\star \in \mathfrak{g}$.

3.3 Examples

In the sequel the stage tree will sometimes be implicitly defined by the transition functions.

For instance, the fourth order commutator-free method [8] is given by

$$\theta_{1,\circ} = \frac{1}{2}F_\circ \tag{41a}$$
$$\theta_{2,\circ} = \frac{1}{2}F_1 \tag{41b}$$
$$\theta_{3,1} = -\frac{1}{2}F_\circ + F_2 \tag{41c}$$
$$\theta_{4',\circ} = \frac{1}{4}F_3 + \frac{1}{6}(F_1 + F_2) - \frac{1}{12}F_\circ \tag{41d}$$
$$\theta_{\bullet,4'} = \frac{1}{4}F_\circ + \frac{1}{6}(F_1 + F_2) - \frac{1}{12}F_3 \tag{41e}$$

It means that we have the vertices $\circ$, 1, 2, 3, $4'$, and $\bullet$. Each equation gives an edge, so there is an edge between $\circ$ and 1, between $\circ$ and 2, etc. The stage tree thus takes the following form



(42)

As mentioned earlier, we also indicate the intermediate vertices, i.e., the vertices for which the component in the F vector is not computed. For instance in the current example, the values $F_{4'}$ and $F_\bullet$ are never needed. Note that the vertex numbering is in general arbitrary, but in this case it indicates the order of the equation for which the method is explicit.

3.3.1 Runge–Kutta methods in $\mathbf{R}^d$

A regular Runge–Kutta method on $\mathbf{R}^d$ with Butcher tableau A, b [11, § II.1.1] has the following skeleton

- The motion map is $\Psi = \exp$
- The transition functions are

$$\theta_{i,\circ} \coloneqq \sum_{j=1}^{s} a_{i,j} F_j \qquad 1 \le i \le s \tag{43a}$$

$$\theta_{\bullet,\circ} \coloneqq \sum_{j=1}^{s} b_j F_j \tag{43b}$$

3.3.2 RKMK4

The fourth order RKMK method [16] may be encoded as follows

$$\theta_{1,\circ} = \frac{1}{2} F_\circ \tag{44a}$$

$$\theta_{2,\circ} = \frac{1}{2} F_1 + \frac{1}{24} [F_\circ, F_1] \tag{44b}$$

$$\theta_{3,\circ} = F_2 + \frac{1}{6} [F_\circ, F_2] \tag{44c}$$

$$\theta_{\bullet,\circ} = V + [I, V] \tag{44d}$$

where

$$V = \frac{1}{6}F_\circ + \frac{1}{3}(F_1 + F_2) + \frac{1}{6}F_3 \tag{44e}$$

$$I = \frac{1}{8}F_\circ + \frac{1}{12}(F_1 + F_2) - \frac{1}{24}F_3 \tag{44f}$$

3.3.3 CG3

The third order Crouch–Grossman method [19] is given by the following transition functions:

$$\theta_{1,\circ} = \frac{3}{4}F_\circ \tag{45a}$$

$$\theta_{2,2'} = \frac{119}{216}F_1 \qquad \theta_{2',\circ} = \frac{17}{208}F_\circ \tag{45b}$$

$$\theta_{\bullet,3'} = \frac{13}{51}F_2 \qquad \theta_{3',3''} = -\frac{2}{3}F_1 \qquad \theta_{3'',\circ} = \frac{24}{17}F_\circ \tag{45c}$$

3.3.4 Symmetric Gauss of order four

The symmetric Gauss method [25] can also be encoded in this way.



(46)

The auxiliary variables $\overline{F}_1$ and $\overline{F}_2$ are determined implicitly by the equation

$$\overline{F}_1 = F_1 + \frac{\sqrt{3}}{4}[\overline{F}_2, F_1] \tag{47a}$$

$$\overline{F}_2 = F_2 - \frac{\sqrt{3}}{4}[\overline{F}_1, F_2] \tag{47b}$$

3.4 Order Zero and Exact Solution

For a given $\xi \in \mathfrak{g}$, we define the ***constant isotropy choice*** $\nu \in \mathcal{C}^\infty(\mathcal{M}, \mathfrak{g})$ as

$$\nu \colon x \mapsto \xi, \qquad x \in \mathcal{M}. \tag{48}$$

We use the abuse of notation

$$\nu = \xi \tag{49}$$

in this case.

Definition 3.1. We say that a skeleton Σ has ***order zero*** if it gives the exact solution for constant isotropy choices, i.e., using the notation (49),

$$\Sigma(\xi) = \exp(\xi) \qquad \xi \in \mathfrak{g}. \tag{50}$$

All the skeletons in § 3.3 have order zero, so one obtains the exact solution with the motion map exp, and using the tautological isotropy map (17) in any of those skeletons.

3.5 Equivariance of Skeletons

The following result shows that in practice it suffices to check the equivariance of the motion map Ψ and the transition functions $\theta_{i,j}$ to obtain an equivariant skeleton, and thus an equivariant method when used with an equivariant isotropy map.

Proposition 3.2. *We have the following "trickling down" results on equivariance:*

(i) *if Ψ and $\theta_{i,j}$ are equivariant, then so are $\Psi_{i,j} = \Psi \circ \theta_{i,j}$*

(ii) *if $\Psi_{i,j}$ are equivariant, then so is the skeleton Σ*

(iii) *if the skeleton Σ and the isotropy map ϖ are equivariant, then so is the method $\Phi = \Sigma \circ \varpi$.*

Proof. We give a proof of item (ii), the other two statement being clear. We simply show that if $x_1 = \Sigma(\nu)(x_0)$ is a solution with the stages X_i and frozen vector fields F_i, then we have $g \cdot x_1 = \Sigma(g \cdot \nu)(g \cdot x_0)$.

$$\begin{aligned} g \cdot X_i &= g \cdot (\Psi_{i,j}(F) \cdot X_j) && \text{(51a)} \\ &= g \cdot (\Psi_{i,j}(F)\, g^{-1}\, g \cdot X_j) && \text{(51b)} \\ &= \Psi_{i,j}(g \cdot F) \cdot g \cdot X_j && \text{(51c)} \end{aligned}$$

□

Note that in all practical examples, the motion maps are of the form $\Psi_{i,j} = \Psi \circ \theta_{i,j}$. The motion map Ψ is the exponential, or Cayley map, both of which are equivariant. The transition functions $\theta_{i,j}$ are Lie-algebra morphisms, as they are linear combinations of commutators.

This gives the following result:

Proposition 3.3. *All the RKMK, Crouch–Grossman and commutator-free skeletons are equivariant.*

3.6 Locality

We show the relation between locality of the skeleton and of the isotropy map. First, we give a practical way of checking that a skeleton is local.

Definition 3.4. We say that the motion map Ψ is local if

$$\Psi(0) = \mathsf{Id} \tag{52}$$

We say that the motion maps $\Psi_{i,j}$ are local if

$$\Psi_{i,j}(0) = \mathsf{Id} \tag{53}$$

We say that the transition functions $\theta_{i,j}$ are local if

$$\theta_{i,j}(0) = 0 \tag{54}$$

Proposition 3.5. *We have the following "trickle down" result on locality.*

(i) *if the motion map Ψ and the transition functions $\theta_{i,j}$ are local, then so are the movement maps $\Psi_{i,j} = \Psi \circ \theta_{i,j}$*

(ii) *if the motion maps $\Psi_{i,j}$ are local, then so is the correspondent skeleton Σ.*

(iii) *if the skeleton Σ and the isotropy map ϖ are local, then so is the method $\Phi = \Sigma \circ \varpi$.*

Proof. The only non trivial statement is the last one. Suppose that $f(x) = 0$ in a neighbourhood of the initial condition x_0. By definition of locality, we then have $\nu(f)(x_0) = 0$. It is then easy to check that taking all the stages X_i equal to x_0 provides a solution, and in particular, we obtain $x_1 = x_0$, which finishes the proof. □

As we already noticed in § 2.3, the tautological isotropy map is *not* local. We saw however in § 3.4 that the corresponding "method" is the exact solution As the exact solution is a local method, the locality of the isotropy map ϖ is *not* necessary.

4 Connections and Reductivity

We study the isotropy maps that are one-forms, i.e., that only depend on the value of the vector field at a given point $x \in \mathcal{M}$, and not on its derivatives. These forms can thus be regarded as "zero order" isotropy maps. The main results of this section are summed up in Figure 4.

4.1 Connections

We define connections as equivariant $\mathfrak{g}$-valued one-forms fulfilling a consistency condition.



Figure 4: There is an affine bijection between the affine space of connections, of invariant connections and of reductive structures. All these spaces are isomorphic to the affine space $\mathcal{L}_{\mathsf{H}}(\mathfrak{g}/\mathfrak{h} \leftarrow \mathfrak{g})$ of H-invariant linear sections from $\mathfrak{g}/\mathfrak{h}$ to $\mathfrak{g}$.

Definition 4.1. We define an ***isotropy form*** as a $\mathfrak{g}$-valued one-form

$$\omega \in \Omega^1(\mathcal{M}, \mathfrak{g}) \tag{55}$$

satisfying the ***consistency condition***

$$\langle \omega, f \rangle_x \cdot x = f(x). \tag{56}$$

It is immediate that the set of isotropy forms has an *affine structure*, i.e., if ω_1 and ω_2 are isotropy forms, then so is $\theta\omega_1 + (1-\theta)\omega_2$ for any real θ.

Definition 4.2. We say that an isotropy form ω is a ***connection*** if it is *equivariant*:

$$\langle \omega, \mathsf{T} g\, f \rangle_{g\cdot x} = g \cdot \langle \omega, f \rangle_x \qquad g \in \mathsf{G}, \quad x \in \mathcal{M}, \tag{57}$$

where the action of g on $\mathfrak{g}$ is the adjoint action (12).

Note that our definition differs from that of an invariant principal connection, and the relation between the two notions is detailed in § 4.3.

Connections are, in a sense, the simplest possible local equivariant isotropy maps, because they are of *order zero*, i.e., they depend only on the value of a vector field at a point, and not on its higher order derivatives. More general local, equivariant isotropy maps are considered in [24].

Proposition 4.3. *Given an isotropy form $\omega \in \Omega^1(\mathcal{M}, \mathfrak{g})$, define the isotropy map ϖ by $\langle \varpi, f \rangle \coloneqq \imath_f \omega$, that is*

$$\langle \varpi, f \rangle(x) \coloneqq \langle \omega, f \rangle_x. \tag{58}$$

The map ϖ is then a local isotropy map. The isotropy map ϖ is equivariant if and only the isotropy form ω is equivariant, that is, if ω is a connection.

Proof. If f is zero in a neighbourhood of $x \in \mathcal{M}$, then, in particular, $f(x) = 0$, so $\langle \varpi, f \rangle(x) = \langle \omega, f \rangle_x = 0$, which proves the locality property.

We note, using the definitions (58), (10), (57) and (11), that

$$\langle \varpi, g \cdot f \rangle(x) = \langle \omega, g \cdot f \rangle_x = \langle \omega, \mathsf{T} g\, f \rangle_{g^{-1}x} = g \cdot \langle \omega, f \rangle_{g^{-1}x} = \big(g \cdot \langle \varpi, f \rangle\big)(x), \tag{59}$$

which shows that ϖ is equivariant if and only if ω is. □

Note that a connection defines at the origin a linear map

$$\omega_{\mathsf{o}} \in \mathcal{L}(\mathfrak{g}/\mathfrak{h}, \mathfrak{g}) \tag{60}$$

where we use the canonical identification

$$\mathfrak{g}/\mathfrak{h} \equiv \mathsf{T}_{\mathsf{o}}\mathcal{M}. \tag{61}$$

The map ω_{o} is H-equivariant, so

$$\omega_{\mathsf{o}} \in \mathcal{L}_{\mathsf{H}}(\mathfrak{g}/\mathfrak{h}, \mathfrak{g}), \tag{62}$$

where, in general,

$$\mathcal{L}_{\mathsf{H}}(V,W) \coloneqq \{\, \varphi \in \mathcal{L}(V,W) \mid \varphi(g\cdot v) = g\cdot\varphi(v) \,\} \tag{63}$$

denotes the H invariant linear maps from a vector space V to a vector space W, both equipped with a linear G-action.

Now, the infinitesimal action of $\mathfrak{g}$ on $\mathsf{T}_{\mathsf{o}}\mathcal{M} \equiv \mathfrak{g}/\mathfrak{h}$ is just the projection $\xi \to \xi + \mathfrak{h}$. We obtain that the consistency condition (56) becomes

$$\langle \omega_{\mathsf{o}}, f\rangle + \mathfrak{h} = f, \qquad f \in \mathfrak{g}/\mathfrak{h}, \tag{64}$$

so we interpret ω_{o} as a *linear section* of the projection $\mathfrak{g} \to \mathfrak{g}/\mathfrak{h}$. We denote the corresponding affine space

$$\mathcal{L}(\mathfrak{g}/\mathfrak{h} \leftarrow \mathfrak{g}) \coloneqq \big\{\, \omega \in \mathcal{L}(\mathfrak{g}/\mathfrak{h}, \mathfrak{g}) \;\big|\; \langle \omega, f\rangle + \mathfrak{h} = f \quad \forall f \in \mathfrak{g}/\mathfrak{h} \,\big\}. \tag{65}$$

The affine space of H-invariant sections is denoted by

$$\mathcal{L}_{\mathsf{H}}(\mathfrak{g}/\mathfrak{h} \leftarrow \mathfrak{g}) \coloneqq \mathcal{L}(\mathfrak{g}/\mathfrak{h} \leftarrow \mathfrak{g}) \cap \mathcal{L}_{\mathsf{H}}(\mathfrak{g}/\mathfrak{h}, \mathfrak{g}). \tag{66}$$

Proposition 4.4. *The map from the affine space of connections to*

$$\mathcal{L}_{\mathsf{H}}(\mathfrak{g}/\mathfrak{h} \leftarrow \mathfrak{g}) \tag{67}$$

defined by $\omega \mapsto \omega_{\mathsf{o}}$ where ω_{o} is defined in (60), is an affine bijection.

The proof is elementary, and is a simplified version of the extension principle presented in details in [17, § 4.2].

Proof. Pick an element $\nu \in \mathcal{L}_{\mathsf{H}}(\mathfrak{g}/\mathfrak{h} \leftarrow \mathfrak{g})$. It defines an connection defined at the point $x = g\cdot\mathsf{o}$ by

$$\langle \omega, f\rangle_{g\cdot\mathsf{o}} = g\cdot\langle \nu, \mathsf{T}g^{-1} f\rangle \tag{68}$$

One checks that the H-equivariance of ν ensures that this map is well defined, i.e., does not depend on which element $g \in \mathsf{G}$ is chosen such that $x = g\cdot\mathsf{o}$. □

4.2 Reductive decompositions

We proceed to show the relation with the existing concept of reductive decompositions.

Definition 4.5. A ***reductive decomposition*** [18] is a decomposition

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m} \tag{69}$$

such that

$$H\cdot\mathfrak{m} \subset \mathfrak{m}. \tag{70}$$

Note that if H is simply connected, the condition (70) is equivalent to

$$[\mathfrak{h}, \mathfrak{m}] \subset \mathfrak{m}. \tag{71}$$

We identify a complementary subspace to $\mathfrak{h}$ as a section of the projection $\mathfrak{g} \to \mathfrak{g}/\mathfrak{h}$, that is a linear map

$$\mu \colon \mathfrak{g}/\mathfrak{h} \to \mathfrak{g} \tag{72}$$

such that

$$\mu(x) + \mathfrak{h} = x \qquad x \in \mathfrak{g}/\mathfrak{h} \tag{73}$$

We have the following relation between equivariance and reductivity:

Lemma 4.6. *The section μ is reductive if and only if it is H-equivariant, that is*

$$h \cdot \mu(x) = \mu(h \cdot x) \tag{74}$$

Proof. As μ is reductive, we have that

$$h \cdot \mu(x) = \mu(x') \tag{75}$$

for some $x' \in \mathfrak{g}/\mathfrak{h}$. But since μ is a section, we have

$$x' = \mu(x') + \mathfrak{h} = h \cdot \mu(x) + \mathfrak{h} = h \cdot (\mu(x) + \mathfrak{h}) = h \cdot x \tag{76}$$

which proves the claim. □

We thus obtain that reductive structures are equivalent to connections.

Proposition 4.7. *The set of reductive structures is in affine bijection with*

$$\mathcal{L}_{\mathsf{H}}(\mathfrak{g}/\mathfrak{h} \leftarrow \mathfrak{g}) \tag{77}$$

A consequence of Proposition 4.4 and Proposition 4.7 is the following result.

Proposition 4.8. *The affine space of connections and the set of reductive structures are in affine bijection with the affine space $\mathcal{L}_{\mathsf{H}}(\mathfrak{g}/\mathfrak{h} \leftarrow \mathfrak{g})$. The underlying linear space is*

$$\mathcal{L}_{\mathsf{H}}(\mathfrak{g}/\mathfrak{h}, \mathfrak{h}). \tag{78}$$

4.3 Relation with Invariant Principal Connections

Recall that a homogeneous space is a particular instance of a *principal* H*-bundle* [22, 23]. In that context, a *principal connection* is a $\mathfrak{h}$-valued one-form θ which is H-equivariant in the sense that

$$\langle \theta, X\, h \rangle_{g\, h} = h^{-1}\, \langle \theta, X \rangle_g\, h, \qquad h \in \mathsf{H}. \tag{79}$$

To be a principal connection, θ must also satisfy the consistency condition

$$\langle \theta, g\,\xi \rangle_g = \xi, \qquad \xi \in \mathfrak{h}. \tag{80}$$

Finally, such a principal connection θ is *invariant* if

$$\langle \theta, \bar{g}\,X \rangle_{\bar{g}\,g} = \langle \theta, X \rangle_g, \qquad \bar{g}, g \in \mathsf{G}. \tag{81}$$

Proposition 4.9. *Consider a connection ω as defined in Definition 4.1. The $\mathfrak{g}$-valued one-form θ defined by*

$$\langle \theta, X \rangle_g \coloneqq g^{-1}\,X - g^{-1}\,\langle \omega, [X] \rangle_{[g]}\,g, \qquad g \in \mathsf{G}, \quad X \in \mathsf{T}_g\mathsf{G}. \tag{82}$$

takes values in $\mathfrak{h}$ and is an invariant principal connection.

Note that in term of the Maurer–Cartan form ω_- defined in (97), θ is defined as $\langle \theta, X \rangle \coloneqq -\langle \omega_-, X \rangle_g - g^{-1}\,\langle \omega, [X] \rangle_{[g]}\,g$.

Proof. Let us show that θ is indeed $\mathfrak{h}$-valued. Define $\xi \coloneqq X\,g^{-1}$, and $\zeta \coloneqq \langle \omega, [X] \rangle_{[g]}$. Notice that

$$\langle \theta, X \rangle = g^{-1}\,(\xi - \zeta)\,g. \tag{83}$$

The projected vector $[X]$ is $[X] = \xi \cdot [g]$. Now the consistency condition (56) reads $\zeta \cdot [g] = [X] = \xi \cdot [g]$, so, by multiplying on the left by g^{-1} (which corresponds to pushing the vector forward to the identity), we obtain $g^{-1}\,(\xi - \zeta)\,g\,\mathsf{H} = 0$, which shows that

$$g^{-1}\,(\xi - \zeta)\,g \in \mathfrak{h}, \tag{84}$$

which, along with (83) implies $\langle \theta, X \rangle \in \mathfrak{h}$.

Now, choosing $X = g\,\xi$ for $\xi \in \mathfrak{h}$, we obtain $[X] = 0$, so $\langle \theta, X \rangle = g^{-1}\,X = \xi$, and the consistency condition (80) is thus fulfilled.

We obtain the H-equivariance by noticing that $[X\,h] = [X]$, so

$$\langle \theta, X\,h \rangle_{g\,h} = h^{-1}\,g^{-1}\,X\,h - h^{-1}\,g^{-1}\,\langle \omega, [X] \rangle_{[g]}\,g\,h = h^{-1}\,\langle \theta, X \rangle_g\,h. \tag{85}$$

Finally, we obtain the invariance of θ by using that $[\bar{g}\,X] = \mathsf{T}\bar{g}\,[X]$ and the equivariance of ω:

$$\begin{aligned} \langle \theta, \bar{g}\,X \rangle_{\bar{g}\,g} &= g^{-1}\,\bar{g}^{-1}\,\bar{g}\,X - g^{-1}\,\bar{g}^{-1}\,\langle \omega, \mathsf{T}\bar{g}\,[X] \rangle_{\bar{g}[g]}\,\bar{g}\,g && (86)\\ &= g^{-1}\,X - g^{-1}\,\langle \omega, [X] \rangle_g\,g && (87)\\ &= \langle \theta, X \rangle_g. && (88) \end{aligned}$$

□

4.4 Flatness and Symmetry

Since a connection defined in Definition 4.1 can also be regarded as a principal connection (§ 4.3), it has a curvature. It also has a torsion, as it is also an *affine connection* [18]. We will not need the exact formulas for the torsion and curvature[2], and focus on whether the connection is flat (zero curvature), or symmetric (torsion free).

Definition 4.10. We say that a connection (and its corresponding reductive structure $\mathfrak{m}$) is ***symmetric*** (or torsion free) if

$$[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{h}. \tag{89}$$

We say that the connection is ***flat*** (or has zero curvature) if $\mathfrak{m}$ is a Lie subalgebra, that is

$$[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{m}. \tag{90}$$

The connection is thus flat and symmetric if and only if $\mathfrak{m}$ is a trivial Lie algebra, i.e., $[\mathfrak{m}, \mathfrak{m}] = 0$.

In the presence of a symmetric connection, the homogeneous space is called a ***symmetric space***. In particular, Cartan performed a classification of the symmetric spaces with compact isotropy group [5]. We refer to the monograph [12] for further details.

4.5 Existence of Connections

We first give a general theorem of existence of reductive structure (and thus of connections) [15, § X.2].

Proposition 4.11. *If the isotropy group* H *is either compact, or connected and semi-simple, there exists a reductive structure.*

In practice, the results of Proposition 4.11 or Proposition 4.4 are of limited use, because it is preferable to have an explicit expression for the connection. The following result, which proof is left to the reader, is used repeatedly to obtain tractable formulas in the examples of § 5, and to calculate the flatness or symmetry of the connection.

Proposition 4.12. *Suppose that a linear form* $\omega \in \Omega^1(\mathcal{M}, \mathfrak{g})$ *is* G*-equivariant and consistent at the origin, i.e.,* $\langle \omega, f \rangle_{\mathsf{o}} \cdot \mathsf{o} = f(\mathsf{o})$*. Then it is a connection corresponding to the reductive decomposition*

$$\mathfrak{m} = \omega(\mathsf{T}_{\mathsf{o}}\mathcal{M}). \tag{91}$$

[2] The interested reader is referred to [18], or [15, § X.2].

4.6 Descending Property of Skeletons

We show that one can use the lifting property of the connection to obtain an integrator in the group G instead, and that this integrator *descends* to $\mathcal{M}$.

A connection ω allows to *lift* a vector field $f \in \mathfrak{X}(\mathcal{M})$ to a ***lifted vector field*** $f_\omega \in \mathfrak{X}(\mathsf{G})$ by

$$f_\omega(g) \coloneqq \langle \omega, f \rangle_{[g]}\, g. \tag{92}$$

Proposition 4.13. *Assume an equivariant skeleton Σ is defined over a group G. Then the integrator $\Sigma(\omega_+)(f_\omega)$ (where ω_+ is the Maurer–Cartan form defined in (95)) descends to the integrator $\Sigma(\omega)(f)$ on the homogeneous manifold $\mathcal{M}$, i.e., the following diagram commutes:*

$$\begin{array}{ccc} \mathsf{G} & \xrightarrow{\Sigma(\omega_+)(f_\omega)} & \mathsf{G} \\ \downarrow & & \downarrow \\ \mathcal{M} & \xrightarrow{\Sigma(\omega)(f)} & \mathcal{M} \end{array} \tag{93}$$

Proof. On the edge i, j, we obtain the equation $G_i = \Psi_{i,j}(F)\, G_j$, so, as $[G_i] = G_i\, \mathsf{H}$, we have $[G_i] = \Psi_{i,j}(F) \cdot [G_j]$. Notice now that by definition of the lifted equation f_ω, we have $\langle \omega_+, f_\omega \rangle_g = \langle \omega, f \rangle_{[g]}$, so $F_i = \langle \omega_+, f_\omega \rangle_{G_i} = \langle \omega, f \rangle_{[G_i]}$, and we conclude that $[G_\bullet]$ is the image of $[G_\circ]$ by $\Sigma(\omega)(f)$. □

5 Connection Examples

We study connections in homogeneous spaces of interest in numerical analysis. The situation varies a lot. Some homogeneous spaces have no connection at all (Proposition 5.7), some have only one (Proposition 5.3), some have infinitely many (§ 5.1.2). There is also a practical aspect, as the connection is not always available in closed form. We refer to Table 1 for a summary of the study of the examples in this section.

5.1 Lie Group as Homogeneous Spaces

A Lie group can be regarded as a homogeneous space in at least three ways, which we now describe.

5.1.1 Left and Right Actions

The Lie group acts transitively on itself by *left multiplication*:

$$g \cdot g' \coloneqq g\, g' \qquad g, g' \in \mathsf{G}. \tag{94}$$

Manifold	Group	Isotropy	Action	Connection	Sym.	Flat	§
$\mathbf{R}^d$	$\mathsf{H} \ltimes \mathbf{R}^d$	$\mathsf{H} \subset \mathsf{GL}(d)$	$hx + a$	$\begin{bmatrix} 0 & \delta x \\ 0 & 0 \end{bmatrix}$	✓	✓	5.2
Stiefel	$\mathsf{SO}(d)(d)$	$\mathsf{SO}(d)(d-k)$	RQ	$\delta Q\, Q^\intercal$ $-Q\, \delta Q^\intercal$ $-Q\, \delta Q^\intercal\, Q\, Q^\intercal$	×	×	5.3
Sphere		$\mathsf{SO}(d)(d-1)$		$\delta Q\, Q^\intercal - Q\, \delta Q^\intercal$	✓	×	
Isospectral	$\mathsf{SO}(d)(\lvert\kappa\rvert)$	$\mathsf{SO}(d)(\kappa)$	$R\, P\, R^\intercal$	No formula	×	×	5.4
	$\mathsf{SO}(d)(\lvert\kappa\rvert)$	$\mathsf{1}$			×	✓	5.4.2
Grassmannian Projective		$\mathsf{S}\big(\mathsf{O}(k) \times \mathsf{O}(d-k)\big)$ $\mathsf{SO}(d)(d-1)$		$\delta P\, P - P\, \delta P$	✓	×	5.4.1
SPD Matrices	$\mathsf{GL}(d)$	$\mathsf{O}(d)$	$A\, P\, A^\intercal$	No formula	✓	×	5.5
G	G	$\mathsf{1}$	$\bar{g}\, g$	$\omega_+ = \quad \delta g\, g^{-1}$	×	✓	5.1.1
			$g\, \bar{g}^{-1}$	$\omega_- = -g^{-1}\, \delta g$	×	✓	
G	$\mathsf{G} \times \mathsf{G}$	G	$g_1\, g\, g_2^{-1}$	$(\omega_+, \omega_-)/2$	✓	×	5.1.2
				$(0, \omega_-)$	×	✓	
				$(\omega_+, 0)$	×	✓	
Fixed rank matrices	$\mathsf{G} = \mathsf{GL}(m) \times \mathsf{GL}(n)$	$\left(\begin{bmatrix} A_1 & 0 \\ A_2 & C \end{bmatrix}, \begin{bmatrix} B_1 & 0 \\ B_2 & C \end{bmatrix}\right) \in \mathsf{G}$	$A\, M\, B^{-1}$	$\emptyset$	$\emptyset$	$\emptyset$	5.6

Table 1: A summary of the connections (or absence thereof) described in this paper. We refer to the respective section in the last column for the notations used in this table.

The isotropy group in that case is trivial, so the only reductive structure is $\mathfrak{m} = \mathfrak{g}$. The corresponding flat connection is the ***Maurer–Cartan form*** [4]. It is defined by

$$\langle \omega_+, \delta g \rangle_g \coloneqq \delta g \, g^{-1}. \tag{95}$$

Note that this connection is symmetric if and only if G is commutative.

Of course, there is also the corresponding right multiplication action

$$g \cdot g' \coloneqq g' \, g^{-1}, \tag{96}$$

for which the flat connection is

$$\langle \omega_-, \delta g \rangle_g \coloneqq -g^{-1} \, \delta g. \tag{97}$$

5.1.2 Cartan–Schouten Action

There is another way in which G can be a homogeneous manifold [6], [15, § X.2]. The symmetry group is now $\mathsf{G} \times \mathsf{G}$, acting on the manifold $\mathcal{M} = \mathsf{G}$ by

$$(g_1, g_2) \cdot g \coloneqq g_1 \, g \, g_2^{-1}. \tag{98}$$

We choose the origin at the group identity. The isotropy group is then

$$\mathsf{H} = \{ \, (g, g) \mid g \in \mathsf{G} \, \} \equiv \mathsf{G}, \tag{99}$$

with corresponding Lie algebra

$$\mathfrak{h} = \{ \, (\xi, \xi) \mid \xi \in \mathfrak{g} \, \}. \tag{100}$$

Define (with a slight abuse of the notation (95)) the $\mathfrak{g} \times \mathfrak{g}$-valued one-form

$$\langle \omega_+, \delta g \rangle \coloneqq (\delta g \, g^{-1}, 0). \tag{101}$$

The infinitesimal action of $(\xi, \zeta) \in \mathfrak{g} \times \mathfrak{g}$ on $g \in \mathsf{G}$ is given by $\xi \, g - g \, \zeta$, so we obtain consistency at the origin. For equivariance, we check that

$$\begin{aligned} \langle \omega_+, g_1 \, \delta g \, g_2^{-1} \rangle_{g_1 \, g \, g_2^{-1}} &= (g_1 \, \delta g \, g_2^{-1} (g_1 \, g \, g_2^{-1})^{-1}, 0) && (102) \\ &= (g_1 \, \delta g \, g^{-1} \, g_1^{-1}, 0) && (103) \\ &= (g_1, g_2) \cdot (\delta g \, g^{-1}, 0). && (104) \end{aligned}$$

This shows, using Proposition 4.12, that ω_+ is a connection and that $\mathfrak{m}_+ \coloneqq (\mathfrak{g}, 0)$ is a reductive structure. The connection ω_+ is flat, because $\mathfrak{m}_+$ is isomorphic to the Lie algebra $\mathfrak{g}$. It is thus symmetric if and only if G is commutative.

There is also a corresponding connection

$$\langle \omega_-, \delta g \rangle \coloneqq (0, -g^{-1} \, \delta g) \tag{105}$$

associated to the reductive structure $\mathfrak{m}_- = (0, \mathfrak{g})$.

As the set of connections is an affine space, the *mean value*

$$\omega_0 = (\omega_+ + \omega_-)/2 \tag{106}$$

of those two connections is also a connection. It also happens to be a *symmetric* connection. Indeed, the associated subspace is $\mathfrak{m}_0 = \omega_0(\mathsf{T}_\mathsf{o}\mathsf{G}) = \{\, (\xi, -\xi) \mid \xi \in \mathfrak{g} \,\}$, and it is easy to check that $[\mathfrak{m}_0, \mathfrak{m}_0] \subset \mathfrak{h}$, showing that the connection is symmetric. The connection ω_0 is thus flat if and only if G is commutative.

We have seen that, in this case, there are at least two reductive structures (so the dimension of the reductive structure space is at least two), but there may be more [2].

5.2 Affine spaces

We define an ***affine space*** as the manifold $\mathcal{M} \equiv \mathbf{R}^d$ and the group $\mathsf{G} = \mathsf{H} \ltimes \mathbf{R}^d$, where H is a subgroup of $\mathsf{GL}(d)$. We represent an element of G by

$$g = \begin{bmatrix} h & a \\ 0 & 1 \end{bmatrix}, \qquad h \in \mathsf{H}, \quad a \in \mathbf{R}^d, \tag{107}$$

acting on an element

$$\begin{bmatrix} x \\ 1 \end{bmatrix} \tag{108}$$

by matrix multiplication. The action is thus

$$g \cdot x = h\, x + a. \tag{109}$$

We choose the origin at zero, i.e., $\mathsf{o} = 0$, and the isotropy group is then H.

On an affine space, there is always an "obvious" connection given by translations.

Proposition 5.1. *In any affine space $\mathsf{H} \ltimes \mathbf{R}^d$ there is a connection given by*

$$\langle \omega, \delta x \rangle_x = \begin{bmatrix} 0 & \delta x \\ 0 & 0 \end{bmatrix}. \tag{110}$$

We now study under which conditions the connection (110) is the only possible one.

Lemma 5.2. *Consider the semidirect product $\mathsf{G} = \mathsf{H} \ltimes \mathbf{R}^d$, where $\mathsf{H} \subset \mathsf{GL}(d)$. For any linear map*

$$\alpha \colon \mathbf{R}^d \to \mathfrak{h} \tag{111}$$

such that

$$[A, \alpha(x)] = \alpha(A\, x), \qquad A \in \mathfrak{h}, \quad x \in \mathbf{R}^d, \tag{112}$$

the subspace

$$\mathfrak{m} = \left\{ \begin{bmatrix} \alpha(x) & x \\ 0 & 0 \end{bmatrix} \,\middle|\, x \in \mathbf{R}^d \right\} \tag{113}$$

is a reductive structure. Conversely, for any reductive structure, there is a linear map α fulfilling (112), such that $\mathfrak{m}$ is defined by (113).

Proof. An element of $\mathfrak{h}$ can be written as

$$\xi = \begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix}, \qquad A \in \mathfrak{h}. \tag{114}$$

Suppose that $\mathfrak{m}$ is a reductive structure. It is parameterised by $x \in \mathbf{R}^d$ as

$$\zeta = \begin{bmatrix} \alpha(x) & x \\ 0 & 0 \end{bmatrix} \tag{115}$$

for some linear function α. We compute the commutator

$$[\xi, \zeta] = \begin{bmatrix} [A, \alpha(x)] & A\,x \\ 0 & 0 \end{bmatrix}. \tag{116}$$

As $\mathfrak{m}$ is reductive, we obtain $[A, \alpha(x)] = \alpha(A\,x)$. It is straightforward to check the opposite statement. □

We now obtain the following uniqueness result if the isotropy group contains scalings. Note how the presence of scalings in the isotropy group also simplified the study of H-invariant spaces in [17, § 6].

Proposition 5.3. *If the isotropy group contains scalings, then the only reductive structure is $\mathfrak{m} = \mathbf{R}^d$. The only connection is then given by (110).*

Proof. If $\mathfrak{gl}(1) \subset \mathfrak{h}$, it means that $\mathsf{Id} \in \mathfrak{h}$. By using (112) with $A = \mathsf{Id}$, we obtain that $\alpha(x) = 0$, so $\mathfrak{m} = \mathbf{R}^d$. □

5.3 Stiefel Manifolds and spheres

For integers $k \leq n$, the ***Stiefel manifold*** is the set of $n \times k$ matrices Q with orthogonal columns, i.e.,

$$Q^{\mathsf{T}}\,Q = 1. \tag{117}$$

An element $R \in \mathsf{SO}(n)$ acts on Q by

$$R \cdot Q \coloneqq R\,Q. \tag{118}$$

This action is transitive. If we define the origin to be the matrix (in block notation)

$$Q_0 \coloneqq \begin{bmatrix} 0 \\ 1 \end{bmatrix} \tag{119}$$

then the isotropy group consists of the matrices

$$\mathsf{H} = \begin{bmatrix} R & 0 \\ 0 & 1 \end{bmatrix}, \qquad R \in \mathsf{SO}(n-k). \tag{120}$$

So, as is customary, we identify $\mathsf{H} \equiv \mathsf{SO}(n-k)$. The Lie algebra $\mathfrak{g}$ is $\mathfrak{so}(n)$, the space of skew symmetric matrices.

A candidate for a reductive decomposition is the space $\mathfrak{m}$ consisting of matrices of the form

$$\begin{bmatrix} 0 & W \\ -W^\intercal & \Omega \end{bmatrix}, \qquad W \in \mathbf{R}^{(n-k)\times k}, \quad \Omega \in \mathfrak{so}(k). \tag{121}$$

Proposition 5.4. *The space $\mathfrak{m}$ is a reductive structure. The associated connection defined at a vector δQ on a point Q is*

$$\langle \omega, \delta Q \rangle_Q = \delta Q\, Q^\intercal - Q\, \delta Q^\intercal - Q\, \delta Q^\intercal\, Q\, Q^\intercal. \tag{122}$$

The connection is symmetric if and only if $k = 1$, i.e., if the Stiefel manifold is a sphere, in which case the connection simplifies into

$$\langle \omega, \delta Q \rangle_Q = \delta Q\, Q^\intercal - Q\, \delta Q^\intercal. \tag{123}$$

Proof. By differentiating (117) we obtain that a tangent vector δQ at Q satisfies

$$\delta Q^\intercal\, Q + Q^\intercal\, \delta Q = 0. \tag{124}$$

This shows that ω takes its value in $\mathfrak{g} = \mathfrak{so}(n)$.

Recalling the action of $\mathsf{SO}(n)$ on the manifold by left multiplication, the lifted action on tangent vectors is also by left multiplication (as the action is linear). Now, for any matrix $R \in \mathsf{SO}(n)$, we have

$$\langle \omega, R \cdot \delta Q \rangle_{R\cdot Q} = R\, \langle \omega, \delta Q \rangle_Q\, R^\intercal = R \cdot \langle \omega, \delta Q \rangle_Q, \tag{125}$$

which shows the equivariance of ω.

At the origin Q_0, the infinitesimal action of

$$\xi = \begin{bmatrix} \star & W \\ W^\intercal & \Omega \end{bmatrix} \in \mathfrak{g} \tag{126}$$

is

$$\xi \cdot Q_0 = \begin{bmatrix} W \\ \Omega \end{bmatrix}. \tag{127}$$

The connection sends that vector to

$$\begin{bmatrix} 0 & W \\ W^\intercal & \Omega \end{bmatrix}, \tag{128}$$

from which consistency follows.

Finally, the image $\mathfrak{m} = \omega(\mathsf{T}_\circ \mathcal{M})$ consists of matrices of the form (121). We conclude using Proposition 4.12.

Finally, computing an extra diagonal term of the commutator between two elements of $\mathfrak{m}$ shows that it is zero if and only if the lower right term is zero, i.e., $\Omega = 0$. This happens only if $k = 1$. Finally, in this case, the orthogonality condition (124) becomes $\delta Q^\intercal\, Q = 0$, which accounts for the simplification in the connection formula. □

Note that the expression (122) is also obtained in [7], although the authors do not mention the equivariance of that connection.

We also notice that if Q' is orthogonal to both Q and δQ, then the vector field at Q' is zero, that is

$$\langle \omega, \delta Q \rangle_Q \cdot Q' = 0. \tag{129}$$

This is especially intuitive on *spheres*, where it means that the (generalised) axis of rotation is orthogonal to the plane spanned by Q and δQ.

5.4 Isospectral Flows, Lax Pairs, Grassmannians and Projective Spaces

An isospectral manifold is the space of symmetric matrices with a prescribed spectrume [3]. As we shall see, they contain as special cases Grassmannians (and thus projective spaces), as well as principal homogeneous spaces for the rotation group, in which case they are often expressed in the form of a Lax pair.

In order to define the isospectral manifold, we first define a partition of an integer d, representing eigenvalue multiplicities. We define a ***partition*** to be a function $\kappa \colon \mathbf{N} \to \mathbf{N}$, which is *non increasing*, and eventually zero. The ***size*** of the partition is

$$|\kappa| = \sum_{i=0}^{\infty} \kappa_i \tag{130}$$

which is a finite integer.

We define the ***isospectral manifold*** $\mathcal{M}$ associated to a partition κ and a sequence of (necessarily distinct) real eigenvalues λ_i as

$$\mathcal{M} \coloneqq \left\{ P \in \mathbf{R}^{|\kappa| \times |\kappa|} \;\middle|\; P = P^\intercal; \quad P \text{ has eigenvalue } \lambda_i \text{ with multiplicity } \kappa_i \right\}. \tag{131}$$

The ***length*** of the partition indicates the number of distinct eigenvalues. It is defined as the number of non zero elements in κ, i.e. $\#\{\, i \in \mathbf{N} \mid \kappa_i \neq 0 \,\}$.

Examples of partitions are $\kappa = (1)$ of length and size one, $\kappa = (3, 2)$ of size five and length two. The partitions associated to a ***Grassmannian*** have length two, such as (m, n), with two arbitrary positive integers m, n such that $m \geq n$. The ***projective space*** case corresponds to the special case $n = 1$.

To a given partition κ we associate a block matrix representation where the block i, j has size $\kappa_i \times \kappa_j$. Note that the size of the partition gives the size of the matrices, whereas the length of the partition is the number of blocks.

In what follows, we exclude the trivial case of κ having length one, in which case the manifold $\mathcal{M}$ reduces to one point.

The group $\mathsf{SO}(|\kappa|)$ acts on P by adjunction

$$R \cdot P = R\, P\, R^\intercal. \tag{132}$$

We can define the origin Δ to be the diagonal matrix with eigenvalues λ_i with multiplicity κ_i. As symmetric matrices are diagonalisable with orthogonal matrices, we obtain that any matrix in $P \in \mathcal{M}$ can be written as $P = R\,\Delta\,R^{\intercal}$ for some $R \in \mathsf{SO}(|\kappa|)$. This shows that $\mathsf{SO}(|\kappa|)$ acts transitively on $\mathcal{M}$. The isotropy subgroup H is then the set of block matrices of the form

$$\begin{bmatrix} R_1 & & \\ & R_2 & \\ & & \ddots \end{bmatrix} \tag{133}$$

such that $R_i \in \mathsf{O}(\kappa_i)$, and such that the determinant is one. We thus denote the isotropy group by

$$\mathsf{H} = \mathsf{SO}(\kappa) \coloneqq \mathsf{S}\Big(\prod_{i \in \mathbf{N}} \mathsf{O}(\kappa_i)\Big). \tag{134}$$

Note that, since the isotropy group does not depend on the eigenvalues, but only on their multiplicity, this shows that all the isospectral manifolds with the same partition (but possibly different eigenvalues) are isomorphic.

A good candidate for the reductive structure $\mathfrak{m}$ is the subspace of $\mathfrak{so}(|\kappa|)$ which is zero on the block diagonal:

$$\begin{bmatrix} 0 & & \\ & 0 & \\ & & \ddots \end{bmatrix} \in \mathfrak{so}(|\kappa|). \tag{135}$$

Proposition 5.5. *The space $\mathfrak{m}$ is a reductive structure. It further induces*

(i) *a* flat *connection if and only if $\kappa = (1, 1, \ldots, 1)$, in which case it is a principal homogeneous space for $\mathsf{SO}(d)$.*

(ii) *a* symmetric *connection if and only if $\kappa = (k, d-k)$, in which case $\mathcal{M}$ is isomorphic to a Grassmannian. The corresponding connection is given explicitly by*

$$\langle \omega, \delta P \rangle_P \coloneqq \frac{1}{(\Delta\lambda)^2}(\delta P\, P - P\, \delta P), \tag{136}$$

where $\Delta\lambda$ is the difference between the two eigenvalues at hand.

Proof. We denote a generic element of $h \in \mathsf{H}$ which is in the form (133). A typical element of $\mathfrak{m}$ can be written, using the same block conventions,

$$m = \begin{bmatrix} 0 & m_{12} & \ldots \\ m_{21} & 0 & \ldots \\ \vdots & & \end{bmatrix} \tag{137}$$

with $m_{ij} + m_{ji}^{\intercal} = 0$.

1. First, a calculation shows that a product of an element $h \in \mathfrak{h}$ and an element $m \in \mathfrak{m}$ is such that $h\,m \in \mathfrak{m}$, and $m\,h \in \mathfrak{m}$, so, in particular, the commutator $[h, m] \in \mathfrak{m}$, which shows that $\mathfrak{m}$ is a reductive structure.

2. The first extra diagonal term, of index $(2, 1)$ in the product $m\,\bar{m}$ of two elements $m, \bar{m} \in \mathfrak{m}$ is

$$\underbrace{m_{11}}_{=0}\,\bar{m}_{12} + m_{12}\,\underbrace{\bar{m}_{22}}_{=0} + m_{13}\,\bar{m}_{32} + \cdots \tag{138}$$

   This shows that if the length of the partition κ is two, then the product is zero at the extra diagonal, and in particular, the commutator is as well, so the connection is symmetric. If the length is greater than two, the commutator contains at least the term

$$m_{13}\,\bar{m}_{32} - \bar{m}_{13}\,m_{32}, \tag{139}$$

   so it is always possible to choose, for instance the blocks m_{13} and $\bar{m}_{32}$, such that this term does not vanishes. This shows that the connection is not symmetric.

3. An element in the block diagonal of the commutator $[m, \bar{m}]$ can be written as

$$\sum_k -m_{ik}\,\bar{m}_{ik}^{\intercal} + \bar{m}_{ik}\,m_{ik}^{\intercal} \tag{140}$$

   If all the blocks have size one, then this vanishes, showing that if all the eigenvalues have multiplicity one, the connection is flat. Note that, in that case, the isotropy group is trivial, which shows that $\mathcal{M}$ is a principal homogeneous space for $\mathsf{G} = \mathsf{SO}(|\kappa|)$. If a block of index k has size greater than one, then one can choose m_{ik} and $\bar{m}_{ik}$ so that $-m_{ik}\,\bar{m}_{ik}^{\intercal} + \bar{m}_{ik}\,m_{ik}^{\intercal} \neq 0$.

4. The equivariance property of (136) is straightforward to check. It now suffices to check consistency at the origin. Recall that we are considering the case

$$\Delta = \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} \tag{141}$$

   with the same block-matrix conventions as before, i.e., with respect to the partition $(k, d-k)$. From (132), we obtain that the infinitesimal action of $\xi \in \mathfrak{so}(d)$ at a point P is given by

$$\xi \cdot P = \xi\,P - P\,\xi. \tag{142}$$

   A direct calculation shows that the commutator of a matrix A with block entries a_{ij} and Δ has block coefficients $[A, \Delta]_{ij} = a_{ij}(\lambda_j - \lambda_i)$. Now, an element of $\mathfrak{m}$ takes the form

$$m = \begin{bmatrix} 0 & m_{12} \\ -m_{12}^{\intercal} & 0 \end{bmatrix}, \tag{143}$$

so $[m, \Delta]$ is a symmetric block matrix

$$\delta\Delta = \begin{bmatrix} 0 & m_{12}(\lambda_2 - \lambda_1) \\ m_{12}^\intercal(\lambda_2 - \lambda_1) & 0 \end{bmatrix}. \tag{144}$$

Applying the commutator $[\delta\Delta, \Delta]$ gives the matrix with element $m_{ij}(\lambda_2 - \lambda_1)^2$. We thus obtain $[\delta\Delta, \Delta] = (\lambda_2 - \lambda_1)^2 m$, which finishes the proof.

□

5.4.1 The symmetric case: Grassmannians

Note that Grassmann manifolds are generally defined as the set of orthogonal projectors on subspaces of dimension k [11, § IV.9.2]. It means that the eigenvalues of the corresponding isospectral manifold are zero and one, so (136) simplifies into

$$\langle \omega, \delta P \rangle_P = \delta P\, P - P\, \delta P. \tag{145}$$

From an element $Q \in \mathbf{R}^{n\times k}$ in a Stiefel manifold (see § 5.3), one obtains an element P in its corresponding Grassmann manifold by

$$P = Q\, Q^\intercal. \tag{146}$$

Grassman manifolds can thus be regarded as the base manifold of a Stiefel manifold considered as a principal bundle. This might be exploited by constructing descending integrators on the Stiefel manifold, instead of the Grassmann manifolds, as we discuss in the conclusion.

5.4.2 The flat case: integrable Lax pairs

The differential equation on an isospectral manifold is often given directly by the infinitesimal action of $\mathfrak{so}(d)$, in which case it is called a *Lax pair*. Indeed, from (142), we have

$$P' = \xi(P) \cdot P = \xi(P)\, P - P\, \xi(P). \tag{147}$$

A particularly interesting case of Lax Pair system is the *integrable case*, when all the eigenvalues are distinct, as for instance, the Toda lattice [11, § X.1.5]. As we saw in Proposition 5.5, this corresponds to the flat case. The isospectral manifold is then a principal homogeneous space, so there is no ambiguity in the isotropy, and the only possibility is thus

$$\langle \omega, \delta P \rangle = \xi(P), \tag{148}$$

so the Lax pair already gives the solution of the connection problem. This is essentially the method considered in [3], and we see now that this method is equivariant. We refer to that aforementioned paper and to [11, § X.1.5] for further insights into Lax pairs and integrable systems.

5.5 Polar Decompositions

There is a homogeneous manifold naturally associated with the polar decomposition of an invertible matrix in an orthogonal and a positive-definite matrix [10, § 3.2]. The group is

$$\mathsf{G} = \mathsf{GL}(d) \tag{149}$$

and the manifold $\mathcal{M}$ is the set of ***symmetric positive definite matrices***:

$$\mathcal{M} = \left\{ P \in \mathbf{R}^{d\times d} \;\middle|\; P = P^\intercal \qquad x^\intercal P\, x > 0 \quad \forall x \in \mathbf{R}^d \setminus 0 \right\}. \tag{150}$$

The action of $A \in \mathsf{GL}(d)$ on $P \in \mathcal{M}$ is given by

$$A \cdot P \coloneqq A\, P\, A^\intercal. \tag{151}$$

It is a transitive action as any positive definite matrix P can be written as $P = A\, A^\intercal$ for some matrix $A \in \mathsf{GL}(d)$. We choose the origin

$$P_0 \coloneqq 1, \tag{152}$$

which gives the isotropy group

$$\mathsf{H} = \mathsf{O}(d). \tag{153}$$

We have the following connection.

Proposition 5.6. *The space of symmetric matrices* $\mathfrak{m} \coloneqq \{\, \xi \in \mathfrak{gl}(d) \mid \xi = \xi^\intercal \,\}$ *is a symmetric reductive structure. The corresponding symmetric connection is defined implicitly by the equations*

$$P\, \langle \omega, \delta P \rangle_P + \langle \omega, \delta P \rangle_P\, P = \delta P \tag{154a}$$

$$\xi = \xi^\intercal. \tag{154b}$$

Proof. The commutator of an antisymmetric matrix and a symmetric matrix is a symmetric matrix, so $[\mathfrak{h}, \mathfrak{m}] \subset \mathfrak{m}$. Moreover, we have $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$, so $\mathfrak{m}$ is a reductive structure. The commutator of two symmetric matrices is an skew symmetric matrix, i.e., $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{h}$ and the connection is thus symmetric. As the infinitesimal action of $\xi \in \mathfrak{g}$ on $P \in \mathcal{M}$ is defined by $\xi \cdot P = \xi\, P + P\, \xi^\intercal$, and as $\xi^\intercal = \xi$ for $\xi \in \mathfrak{m}$, we obtain that (154a) has a unique solution in $\mathfrak{m}$. □

5.6 Matrices of fixed rank

The space of matrices of fixed rank is considered in [11, § IV.9.3] and plays a fundamental role in low-rank approximations, and model reduction. We show how this space has a natural structure of homogeneous space, and that it lacks reductive structures, and thus, connections.

For integers m, n, and k, we define the manifold $\mathcal{M}$ of $m \times n$ matrices of rank k. An element $(A, B) \in \mathsf{GL}(m) \times \mathsf{GL}(n)$ acts on such a matrix M by

$$(A, B) \cdot M = A\, M\, B^{-1}, \tag{155}$$

and the action is transitive on $\mathcal{M}$.

Let us choose the origin at the $m \times n$ matrix M_0 of rank k defined by

$$M_0 \coloneqq \begin{bmatrix} 0 & 0 \\ 0 & \mathsf{1} \end{bmatrix}, \tag{156}$$

where $\mathsf{1}$ denotes here the identity matrix of size k. We assume the relevant block decomposition for the matrices A, B and M in the remaining of this section.

A calculation shows that the isotropy group consists of pairs of matrices of the form

$$\left(\begin{bmatrix} A_1 & 0 \\ A_2 & C \end{bmatrix}, \begin{bmatrix} B_1 & 0 \\ B_2 & C \end{bmatrix} \right) \in \mathsf{GL}(m) \times \mathsf{GL}(n). \tag{157}$$

Proposition 5.7. *The homogeneous space of $m \times n$ matrices of rank k has a connection if and only if $m = n = k$, in which case it is the Cartan–Schouten manifold (see § 5.1.2) for $\mathsf{GL}(k)$.*

Proof. First, we notice that if $m = n = k$, then the rank k matrices are simply the invertible matrices of size k, and the action (155) is the Cartan–Schouten action (98).

We now assume that there exists a reductive structure, and proceed to show that it implies $m = n = k$.

A reductive structure $\mathfrak{m} \subset \mathfrak{g}$ is parameterised by matrices α, β, γ, and linear maps $\overline{a_1}$, $\overline{a_2}$, $\overline{b_1}$, $\overline{b_2}$ and $\overline{c}$, all depending linearly on α, β, γ, such that an element $m \in \mathfrak{m}$ takes the form

$$m = \left(\begin{bmatrix} \overline{a_1} & \alpha \\ \overline{a_2} & \overline{c} - \gamma \end{bmatrix}, \begin{bmatrix} \overline{b_1} & \beta \\ \overline{b_2} & \overline{c} + \gamma \end{bmatrix} \right). \tag{158}$$

We denote the infinitesimal adjoint action of an element $\xi \in \mathfrak{h}$ by

$$\xi \cdot m \coloneqq [\xi, m] = \xi\, m - m\, \xi. \tag{159}$$

1. Choose

$$\xi = \left(\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \right) \in \mathfrak{h}. \tag{160}$$

We obtain that

$$\xi \cdot m = \left(\begin{bmatrix} 0 & \alpha \\ -\overline{a_2} & 0 \end{bmatrix}, \begin{bmatrix} 0 & \beta \\ -\overline{b_2} & 0 \end{bmatrix} \right), \tag{161}$$

so we obtain

$$\overline{a_2}(\alpha, \beta, 0) = -\overline{a_2}(\alpha, \beta, \gamma). \tag{162}$$

By choosing $\gamma = 0$, we first obtain $\overline{a_2}(\alpha, \beta, 0) = 0$, and thus also $\overline{a_2}(\alpha, \beta, \gamma) = 0$ for all values of the parameters α, β, γ.

2. Choose now an arbitrary element

$$m = \left(\begin{bmatrix} \overline{a_1} & \alpha \\ 0 & \tilde{c} \end{bmatrix}, \star \right) \in \mathfrak{m}, \tag{163}$$

where $\star$ denotes here and in the sequel an arbitrary element. Choose an element

$$\xi = \left(\begin{bmatrix} 0 & 0 \\ A & 0 \end{bmatrix}, \star \right) \in \mathfrak{h}. \tag{164}$$

We obtain

$$\xi \cdot m = \left(\begin{bmatrix} -\alpha\, A & 0 \\ A\, \overline{a_1} - \tilde{c}\, A & A\, \alpha \end{bmatrix}, \star \right), \tag{165}$$

so we obtain the condition

$$A\, \overline{a_1} - \tilde{c}\, A = 0. \tag{166}$$

Now, the element $\xi \cdot m$ also belongs to $\mathfrak{m}$, so the condition $\xi \cdot (\xi \cdot m) \in \mathfrak{m}$ gives

$$A\, (-\alpha\, A) - A\, \alpha\, A = 0. \tag{167}$$

3. We choose in particular $A = \alpha^\intercal$, and we get

$$\alpha^\intercal\, \alpha\, \alpha^\intercal = 0. \tag{168}$$

4. Now, the parameter α is arbitrary, so if $m > k$, we can choose it such that either $\alpha\, \alpha^\intercal$ or $\alpha^\intercal\, \alpha$ is the identity matrix, but (168) then gives $\alpha = 0$, which is a contradiction. We conclude that $m = k$. A similar reasoning for β would give $n = k$.

□

The following example of a nonreductive space, mentioned in [15, § X.2],

$$\mathsf{G} = \mathsf{SL}(2, \mathbf{R}), \qquad \mathsf{H} = \left\{ \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix} \;\middle|\; \lambda \in \mathbf{R} \right\}, \tag{169}$$

corresponds to 2×1 matrices which are nonzero (i.e., which have rank one), so it is the case $m = 2$, $n = 1$ and $k = 1$ in Proposition 5.7.

6 Conclusion and Open Problems

The main message of this paper is that equivariant isotropy map allow to construct equivariant homogeneous space integrators from Lie group integrators (skeletons). Moreover, when that equivariant isotropy map is of order zero, then it is equivalent to

a reductive structure or an invariant principal connection, both of which are standard tools in differential geometry [15, § X.2].

We examine the consequences of using a connection with a skeleton.

The first consequence is that some homogeneous spaces currently lack (equivariant) integrators. A fundamental such example, as we showed in § 5.6, is the homogeneous space of matrices of fixed rank.

Another consequence is that, as the connection takes its values in $\mathfrak{m}$, one only need compute exponentials of elements of $\mathfrak{m}$. For instance in the affine case, only translations are needed.

In a similar vein, note that if the motion map is the exponential, one need to know one exponential on each H-orbit since

$$\exp(h \cdot \xi) = h \cdot \exp(\xi) \qquad h \in \mathsf{H} \quad \xi \in \mathfrak{m}. \tag{170}$$

For instance, on a sphere, all the "great circle" are in a sense, the same, as there is only one H-orbit.

Let us mention some open questions to be investigated in future work.

- It is not clear yet whether the isotropy map has to be equivariant for the integrator to be. Isotropy map equivariance is necessary in some specific cases, but we do not expect that result to extend to general cases. Nevertheless, using equivariant isotropy maps is still the simplest way to obtain equivariant homogeneous space integrators, and we are not aware of any concrete example of an equivariant integrator constructed with a non-equivariant isotropy map.

- Homogeneous spaces lacking a connection such as the space of matrix of fixed rank considered in § 5.6 might have higher order connections. Such connections are still local (and thus, computationally tractable), but depend on higher order derivatives of the vector field [24]. We do not know if such a higher order equivariant connection exists for the space of fixed rank matrices.

- As noted in [9], the representation of points on the Grassmannians as $n \times n$ matrices may be unwieldy, so there might be strategies to lift the equation on the Stiefel manifold and integrate there instead. This is the strategy proposed in [11, § IV.9.2], but we do not know if this gives equivariant integrators.

- The retraction methods exposed in [7] on Stiefel manifolds, are not of the same type as those in this paper. We believe that connections may be used to conceive retraction methods of the same type on other homogeneous spaces. We do not know under which conditions retraction methods are equivariant.

- We showed how some homogeneous spaces have a *symmetric* connection. In particular, in the Cartan–Schouten case, one could use the symmetric connection. When used along with skeletons, only the *geodesics* of the connection matter, so all the connections studied in § 5.1.2 will give the same methods, but there might be a way to exploit the symmetric connection.

- In Proposition 5.3 we show that the connection is unique, under some assumptions. We don't know of any such results for the other homogeneous spaces of interest, in particular, those presented in § 5.
- Finally, we mention other homogeneous spaces which are used in physics, such as the Galilean and Poincaré groups, and hyperbolic spaces such as the Poincaré half space $\mathsf{SL}(d)/\mathsf{SO}(d)$, the de Sitter space, or the Lagrangian Grassmannian $\mathsf{U}(d)/\mathsf{O}(d)$ [21], the flag manifolds [14, § 2.5], not to mention the complex and quaternionic variants of the Stiefel and Grassmann manifolds. Note that, even if there may be theoretical results of existence of connection, in particular when the isotropy group is compact, as we saw in Proposition 4.11, it does not mean that there is an actual, practically computable connection formula. Such examples are given by the isospectral manifolds § 5.4 and the symmetric positive definite matrices § 5.5, for which the connection exists, but may be difficult to compute.

Acknowledgements

This research was supported by the Spade Ace Project and by the J.C. Kempe memorial fund.

References

[1] R. Abraham, J. E. Marsden, and T. Ratiu. *Manifolds, Tensor Analysis, and Applications*, volume 75 of *Applied Mathematical Sciences*. Springer-Verlag, New York, second edition, 1988. ISBN 0-387-96790-7.

[2] R.E. Beck. Connections on semisimple Lie groups. *Trans. Am. Math. Soc.*, 164: 453–460, 1972. ISSN 0002-9947; 1088-6850/e. doi: 10.2307/1995989.

[3] M. P. Calvo, A. Iserles, and A. Zanna. Numerical solution of isospectral flows. *Math. Comput.*, 66(220):1461–1486, 1997. ISSN 0025-5718; 1088-6842/e. doi: 10.1090/S0025-5718-97-00902-2.

[4] É. Cartan. La structure des groupes de transformations continus et la théorie du trièdre mobile. *Bull. Sci. Math., II. Sér.*, 34:250–283, 1910. ISSN 0007-4497.

[5] É. Cartan. Sur une classe remarquable d'espaces de Riemann. I. *Bull. Soc. Math. Fr.*, 54:214–264, 1926. ISSN 0037-9484.

[6] É. Cartan and J. Schouten. On the geometry of the group-manifold of simple and semi-simple groups. *Nerl. Akad. Wensch. Proc. Ser. A*, 29:803–815, 1926.

[7] E. Celledoni and B. Owren. On the implementation of Lie group methods on the Stiefel manifold. *Numer. Algorithms*, 32(2-4):163–183, 2003. ISSN 1017-1398; 1572-9265/e. doi: 10.1023/A:1024079724094.

[8] E. Celledoni, A. Marthinsen, and B. Owren. Commutator-free lie group methods. *Future Gener. Comput. Syst.*, 19(3):341–352, April 2003. ISSN 0167-739X. doi: 10.1016/S0167-739X(02)00161-9.

[9] A. Edelman, T. A. Arias, and S. T. Smith. The geometry of algorithms with orthogonality constraints. *SIAM J. Matrix Anal. Appl.*, 20(2):303–353, 1998. ISSN 0895-4798; 1095-7162/e. doi: 10.1137/S0895479895290954.

[10] J. Gallier. Notes on differential geometry and lie groups. 2014. URL http://www.cis.upenn.edu/~jean/gbooks/manif.html.

[11] E. Hairer, C. Lubich, and G. Wanner. *Geometric numerical integration: structure-preserving algorithms for ordinary differential equations*. Springer series in computational mathematics. Springer, 2006. ISBN 9783540306634.

[12] S. Helgason. *Differential Geometry and Symmetric Spaces*. Pure and applied mathematics. American Mathematical Society, 1962. ISBN 9780821873175.

[13] A. Iserles, H. Z. Munthe-Kaas, S. P. Nørsett, and A. Zanna. Lie-group methods. In *Acta Numerica. Vol. 9*, pages 215–365. Cambridge: Cambridge University Press, 2000. ISBN 0-521-78037-3/hbk.

[14] A. Kirillov. *An introduction to Lie groups and Lie algebras*. Cambridge: Cambridge University Press, 2008. ISBN 978-0-521-88969-8/hbk.

[15] S. Kobayashi and K. Nomizu. *Foundations of differential geometry. Vol. II*. Wiley, 1969. ISBN 9780470496480.

[16] H. Munthe-Kaas. Runge-Kutta methods on Lie groups. *BIT*, 38(1):92–111, 1998. ISSN 0006-3835; 1572-9125/e. doi: 10.1007/BF02510919.

[17] H. Z. Munthe-Kaas and O. Verdier. Equivariant series, 2013. URL http://arxiv.org/abs/1308.5824.

[18] K. Nomizu. Invariant affine connections on homogeneous spaces. *Am. J. Math.*, 76:33–65, 1954. ISSN 0002-9327; 1080-6377/e. doi: 10.2307/2372398.

[19] B. Owren and A. Marthinsen. Runge-Kutta methods adapted to manifolds and based on rigid frames. *BIT*, 39(1):116–142, 1999. ISSN 0006-3835; 1572-9125/e. doi: 10.1023/A:1022325426017.

[20] X. Pennec, P. Fillard, and N. Ayache. A riemannian framework for tensor computing. *International Journal of Computer Vision*, 66(1):41–66, 2006.

[21] P. Piccione and D. V. Tausk. *A student's guide to symplectic spaces, Grassmannians and Maslov index*. Rio de Janeiro: Instituto Nacional de Matemática Pura e Aplicada (IMPA), 2008. ISBN 978-85-244-0283-8/pbk.

[22] R.W. Sharpe. *Differential geometry: Cartan's generalization of Klein's Erlangen program*. Graduate texts in mathematics. Springer, 1997. ISBN 9780387947327.

[23] S. Sternberg. Lectures on differential geometry. Prentice-Hall Mathematics Series. Englewood Cliffs: Prentice-Hall, Inc. xi, 390 pp. (1964)., 1964.

[24] O. Verdier. Integrators on homogeneous spaces: local isotropy maps. in preparation, 2014.

[25] A. Zanna, K. Engø, and H. Z. Munthe-Kaas. Adjoint and selfadjoint Lie-group methods. *BIT*, 41(2):395–421, 2001. ISSN 0006-3835; 1572-9125/e. doi: 10.1023/A:1021950708869.